The Impact of Indoor Training Program Due to covid19 quarantine, on Some Physiological, Physical variables and performance efficiency in Elite Wrestlers

¹ ^a MARWA FROUQ GHAZY, ² ^b MOHAMED SAAD GHORABA

¹ Department of Sport Health Sciences, Tanta University, Egypt

^a d.marwa.ghazy@phed.tanta.edu.eg, orcid.org/0000-0002-6109-3939

² Department of Combat and individual sports, Tanta University, Egypt

^b mohamed.ghoraba@phed.tanta.edu.eg

Abstract

Egypt's COVID-19 co-existence plan insisted on social distancing and keeping closure of physical and sport entities. WHO illustrated the importance of training at home with at least 150 minutes of moderate load exercises or 75 minutes of high load exercises in order to reduce the indoor related health problems such as; anxiety, stress, immunity deficiency, so the study aimed to investigate the Impact of Indoor Training Program Due to covid19 quarantine, on Some Physiological, Physical, and Technical Variables in Elite Wrestlers. 8 professional wrestlers with average age 19.3 years, agreed to participate in 12 week of indoor training. Long period change in training regime with short bouts of resistive physical and technical exercises affected the back flexibility with reduction of 20.673%, physiological variables significantly reduced, maximal oxygen consumption "vo2max", breathing frequency "BF" recorded decrease with 8.781%, 6.132% respectively, ratio between carbon dioxide and oxygen "RER" recorded slight nonsignificant increase with 3.316%, and little change in calorie expenditure showing gradual decrease with stay home effect on wrestlers body. Performance efficiency decreased especially in performance endurance measured in time and score with 34.756%, 16.364% respectively, in correlation with the decrease in physiological variables.

Keywords: indoor training, wrestling, efficiency, covid19.

Introduction

COVID-19 pandemic changed the face of the world. Since corona virus had spread nearly all over the world, restricted instruction came out of health ministries and agencies in order to cease spreading of the virus. All aspects of human life was affected with restrictions including; social activities, physical activities and sports, these restrictions resulted closure of sport facilities like sporting clubs, gyms, fitness centers and rehab centers (Daniela et al., 2020). Egypt's COVID-19 co-existence plan also insisted on social distancing and keeping closure of physical and sport entities (State information service, 2020). Stay home was the first goal of all governorates all over the world so, WHO and health monitors began to take a good care of detraining effect on human health and immunity which badly needed for reducing the probability of infection. WHO illustrated the importance of training at home with at least 150 minutes of moderate load exercises or 75 minutes of high load exercises in order to reduce the indoor related health problems such as; anxiety, stress, immunity deficiency and even fear of loss of family and beloved ones (Daniela et al., 2020). Amateurs and professional athletes were the highly affected by stay home restriction, so tele-training, self-training and online personal trainers were and still in action. Detraining was the main concern for trainers where interrupting training has a passive effect on athletes' physiological and physical adaptation that can also lead to abnormality in eating, sleeping and psychological status so (Mujika, and Padilla, 2000; Chen, et al., 2020; Joo, 2018), This was the opportunity for the researchers to carry out this study to identify the capability of tele-training based on interactive training program on elite wrestlers' physiological and physical parameters.

Methods & Materials

Research curriculum: Experimental curriculum used in this study, experimental method with pre-post measurements carried out for one experimental group.

Sample: 8 healthy, professional wrestlers with average age 19.3 years, agreed to participate "with written consent" in this study, wrestlers are members of Egyptian wrestling federation. Table (1), describes the homogenous of the study sample.

Table (1), statistical norms of the study sample in all the study variables

Main variables	unit	Mean	Median	SD	skew	Torsion
Growth variables						
Age	year	19.375	19.500	1.408	-0.564	-0.480
Height	cm	170.250	170.500	3.615	-1.255	-0.221
Weight	kg	75.500	76.000	3.381	-0.796	-0.429
Training experience	year	8.250	8.000	1.282	-1.546	0.475
Physiological variables						
Vo2max	mL/kg/min	47.194	42.415	1.403	2.234	-1.230
BF	L/min	53.000	54.000	3.703	-0.722	0.068
RER	L/min	1.659	1.815	0.366	-1.309	-0.666
Calories	kcal	94.973	96.730	5.450	0.036	-0.930
Physical variables						
Handgrip "right"	kg	35.250	35.000	2.053	0.142	0.743
"Handgrip "left	kg	29.125	29.000	1.808	-0.930	0.336
Back muscle strength	kg	79.625	79.500	3.292	0.700	-0.135
Back muscle flexibility "bridge"	cm	26.000	26.000	3.071	-0.804	0.355
Technical variables						
Performance efficiency "score"	score	20.500	20.000	1.069	-0.831	0.468
Performance efficiency "speed"	sec	6.875	7.000	0.641	0.741	0.068

Coefficient of torsion at significance $0.05=1.474$

Table (1), represents the mean, median, standard deviation "SD" and Coefficient of torsion of the sample members in all study variables, where data values ranged (± 3) bellow Coefficient of torsion.

Experimental work: pretests were taken in frame of preparing to the Egyptian universities' championship in March 16th, to 18th just before university shuts down and closure of all sport facilities, researchers took the opportunity to test the capability of tele-interactive indoor training program in maintaining the athletes' physiological and physical parameters. Study tests were based on pretest before closure; tests were physiological parameters with Ergospirometry. Vo2max (maximum oxygen uptake) recorded in mL/kg/min, BF (breathing frequency) in number, RER (ratio between exerted carbon dioxide and inhaled oxygen) in ratio, and calories expended in the overall test. Physical test were handgrip strength (hand dynamometry), back muscles strength (dynamometry) and back flexibility (with bridge test) in centimeter. Efficiency of performance (score- time) was the technical tests where athletes perform 3 times bridge and throw with time estimating, the shorter time scored, the more efficiency in performing as a performing speed, another test performed by a multiple suplexes with time record, the more suplexes recorded the more efficiency in performing, as a performing endurance. Tests carried out to investigate the influence of indoor training on technical capability of the wrestlers.

Wrestlers experienced tele-interactive training program, consisted of 3 sessions/ week and lasted up to 12 week "April 1st – July 1st ", of indoor training till the gradual reopening of sport facilities in June-2020. Training program was alike the traditional training program of the wrestlers, but

the challenge for this study were the equipment, adequate wrestling mats, open area for full range of motion during performance, face to face wrestling mates and proper recovery. Training program carried out by coaching over zoom cloud meeting app, as it's easy and common in use between youth, Table (2), represents the training program. Post-tests carried out after reopening of sport facilities, between, the 4th and 7th of July.

Table (2), the indoor training program

Week	Session content	Unite Duration	Exercise, (wrestling drills)	Intensity	Rept.	Group	Tools
1 - 4	Warm up	5min.	March in place Muscle stretching	40-50%	1	1	Without
	Main part (wrestling drills)	30min.	arm control "left-right" (cling)	60-75%	20	3	Elastic rubber - pull rally rope- dumbbell- sand bags
			palm, wrist Grip and Penetration by Elastic rubber				
			Snatch drill upright position (obstacle resistance)				
			Bridge skill and get rid of				
	Closure part	5min.	Cool down-Stretch	30%	1	1	Without
5 - 8	Warm up	5min.	Jogging Muscle stretching	40-50%	1	1	Without
	Main part (wrestling drills)	30min.	arm control (cling)	75 - 85%	25	3	Elastic rubber - dumbbells 10k- pull rally rope- kettlebells
			Snatch drill upright position				
			Penetration skill and upper-hunch by Elastic rubber				
			take down skill by Elastic rubber				
			Lifting skill and back-arch by Elastic rubber				
			Snatch drill down position (heavy resistance)				
			palm, wrist Grip and Penetration by Elastic rubber				
			down position (wrenching against obstacle)				
	Closure part	5min.	Jogging	30%	1	1	Without
9 - 12	Warm up	5min.	Jogging Muscle stretching	40-50%	1	1	Without
	Main part (wrestling drills)	30min.	arm control "left-right" (cling)	75-85%	25	3	Elastic rubber - dumbbells 10k- pull rally rope- kettlebells
			palm, wrist Grip and Penetration by Elastic rubber				
			Penetration skill and upper-hunch by Elastic rubber				
			take down skill by Elastic rubber				
			Bridge skill and get rid of				
	Closure part	5min.	Jogging	30%	1	1	Without

Statistical analysis: Statistical analyses were performed using SPSS software, M "means", SD "standard deviation", T-test and Changing percentage were used for treating the collected data of study tests.

Results

Analytic treatment of the study data resulted, the following results, tables 3, 4, represent the study results.

Table (3), t-test and changing percentage of the physiological variables of the study

Physiological variables	Pre-test		Post-test		Mean def.	t-test	Changing percentage
	M	SD±	M	SD±			
Vo2max	47.194	1.403	43.489	1.637	3.705	4.404	8.781
BF	53.000	3.703	49.750	3.412	3.250	3.149	6.132
RER	1.659	0.366	1.714	0.378	0.055	0.459	3.316
Calories	94.973	5.450	96.347	5.884	1.374	0.461	1.446

T value at significance of 0.05=1.895

Table (3), represents significant differences between pre and post-test in vo2max and BF variables, but no significant differences found between pre and post-tests of RER and calorie expended in overall test, were calculated t value was between (0.461:4.404).

Table (4), t-test and changing percentage of the physical and performance efficiency variables of the study

physical and technical variables	Pre-test		Post-test		Mean def.	t-test M	Changing percentage SD±
	M	SD±	M	SD±			
Physical variables							
Handgrip "right"	35.250	2.053	35.975	2.031	0.725	2.339	2.057
"Handgrip "left	29.125	1.808	30.125	1.959	1.000	2.646	3.433
Back muscle strength	79.625	3.292	81.450	3.855	1.825	6.938	2.292
Back muscle flexibility "bridge"	26.000	3.071	31.375	3.420	5.375	11.672	20.673
technical variables							
Performance efficiency "score"	20.500	1.069	13.375	0.916	7.125	7.139	34.756
Performance efficiency "speed"	6.875	0.641	8.000	0.756	1.125	9.000	16.364

T value at significance of 0.05=1.895

Table (4), represents significant differences between pre and post-test in all physical and performance efficiency variables were calculated, t value was between, (2.339, 13.840).

Discussion

Indoor training was a challenging experience of keeping athletes in action, lack of space, partnership cooperation and sport equipment were the first challenge to both athletes and trainers. To investigate the impact of indoor training on athletes' status the researcher chose physiological, physical and technical variables, as representors of athletes' overall wellbeing. Wrestling is a high intensity sport with short interval of performance where wrestlers always attempted to maximize their muscle size and power (Yoon, 2002), although wrestling is characterized with anaerobic system, but lots of studies used vo2max as a detector of wrestlers status (Arslanoğlu, 2015; Ramirez-Velez, et al., 2014), VO2max is considered one of the common indicators of evaluating aerobic capacity and body energy system, VO2max reflects a player's overall fitness level, it is also known as aerobic and oxidative capacity, where it represents the maximum oxygen consumption used during building ATP (Hadžović-Džuvo, et al., 2014), VO2max is characterized by the limits of ones oxygen transport system, so its value does not vary from moment to moment, that gives a true indication of the athletes aerobic capacity (Scribbans, et al., 2016; Dlugosz, et al., 2013).

Physiological variables taken by Ergometer cardiopulmonary test showed significant difference in vo2max reading, The recent study recorded mean of 47.194 mL/kg/min in the pre-test showing moderate status of the study sample where international range of wrestlers' vo2max recorded 41.2 mL/kg/min for United States Freestyle Wrestlers, 45.9 ± 6.6 mL/kg/min for Colombian wrestlers, (Ramirez-Velez, et al., 2014), 49 mL/kg/min for Turkish wrestlers (Arslanoğlu, 2015), and 59.8 mL/kg/min for polish wrestlers (Hübner-Woźniak, et al., 2009). Post-test of vo2max recorded decreased mean with 43.489 mL/kg/min in wrestlers reading after indoor training showing the decrease of aerobic capacity with 2.057% in sake of anaerobic adaptation, that can be Attributed to less of working area, lack of performance endurance where no colleague or opponent to challenge, the attitude of the training program which depended on small resistance with short time endurance, but as a physiological detector, the post reading still in the frame of high trained athletes reading according to the relative studies (Heywood, 2006).

BF (breathing frequency) is the number of breaths per minutes. Adaptive breathing frequency is decreased in trained athletes at rest with 7-8 breaths/minute, where it ranges about 12-20 breaths/minute in untrained individual. This reduction in breathing frequency represents greater respiration efficiency which attributed to adequate training or physical exercises (Gulam, 2016), unlike the measured BF during exercise where breathing frequency rises related to the high efficiency of athletes' respiration (Nicolò, et al., 2017). Recent reading of BF showed a little reduction with 6.132%, related to vo2max reduction showing minor retreat of respiration efficiency caused by the less endurance enclosed in the indoor training program.

RER is the ratio between the amount of exerted carbon dioxide (CO₂) exhaled and the inhaled oxygen (O₂) showing the energy expenditure during the performance (Ramos-Jiménez, et al., 2008). RER is related to the high training intensity that could determine the anaerobic threshold of the athletes (Bellar, and Judge, 2012), matching the wrestling energy regime. RER value is considered a proper indicator of overall physical fitness (Bearden, et al., 2004). Recent RER reading recorded insignificant increase with 3.316% for the post-test where related studies emphasized unchanging values of RER in case of changing the training intensity (Houmard, et al., 1992), RER reduction signaling a drop in aerobic metabolism in sake of carbohydrate consumption matching the results of related studies (Bellar, and Judge, 2012; Rietjens, 2001), recent reading also related to decrease in vo2max and BF collected from post-test of the sample wrestlers showing the physiological changing in anaerobic capacity and fatigue threshold.

Insignificant increase in calorie expended in the post-test with 1.446%, reflecting the change in energy expenditure and fast fatigue threshold of the wrestlers after the reduction of the training bout "indoor training" matching the results of related studies (Madsen, 1993; Mujika, and Padilla, 2001).

The training program in this study cared avoiding the detraining condition that may result out of "stay home" restriction, although the detraining influence the athletes after a long period of cutting the training off (Mujika, and Padilla, 2001), but studies pointed that a less amount of training may help in reducing the impact of detraining effect (MacDougall, and Sale, n.d.), that was the priority of this study, the researcher tried to preserve the physical and technical capabilities as well, so the training program depended on short bouts of specific physical exercises, the physical pre/post-tests of the wrestlers were significantly changed as little increase with 2.057% and 3.433%, in muscular strength represented in hand grip test for right and left handgrip respectively, where

hand grip strength is considered a detector of overall body strength and successful wrestling performance (García-Pallarés, et al., 2011; Iermakov, et al., 2016), the recorded increased in right and left hand grip test is due to the focus of the training program on single joint action strength exercises as it's a successful method of increasing the overall body strength (Gentil, et al., 2015), and because of the ease of its performance and the less equipment needed, that was convenient with home training. Back muscle strength test as well is considered a successful method of testing wrestlers strength capability (Guilhem, et al., 2014), where relative studies emphasized on its crucial impact in wrestling performance (García-Pallarés, et al., 2011; Kraemer, et al., 2001), recent study resulted significant increase in back muscle reading in post-test with 2.292%, which also directs the influence of single joint muscle strengthening in wrestlers as performed in the study program matching the related study which showed the slow effect of reducing the exercising bouts on isometric strength of back muscles (de França, et al., 2015; Tucci, et al., 1992). On the contrary, back mobility test recorded significant reduction with 20.673% as the single joint resistance exercise resulted in a strong but stiff body in contrast with resistive wide range of motion resulted in technical performance with actual opponent weight, then the applied exercises of single joint increased the muscles strength but lowered the mobility of back muscle which is compatible with related studies (Basar, et al., 2014; Chaabene, et al., 2017) unlike the findings of other related studies which demonstrated a little increase in back flexibility out of muscle strengthening only (Leite, et al., 2017; Saraiva, et al., 2014). Technical variables recorded significant reduction with 34.756% for performance efficiency in score, and 16.364% for performance efficiency in time, which represent the performance endurance. This reduction was related to previous decrease in physiological and physical variables indicating the integrating adaptation of athlete's body (Swärd, 1990).

Conclusions

Wrestling performance influenced by the change in training regime due to "stay home" covid19 restrictions. Although athletes maintained their muscle strength, but the long period change in training regime with short bouts of resistive physical and technical exercises affected the back flexibility and physiological variables with reduction in vo2max, BF, RER, and little change in calorie expenditure. Performance efficiency decreased especially in performance endurance measured in time and score, in correlation with the physiological variables. Indoor training succeeded in ceasing the influence of detraining, but wasn't sufficient enough to simulate real performing with opponent resistance.

Conflict of interest: the authors declare that this study is self-funded, and there is no conflict of interest.

References

1-Daniela B., Melissa M., Carol P., & Robert V. (2020). The impact of COVID-19 on sport, physical activity and well-being and its effects on social development. UN/DESA Policy Brief.;73:1-5

- 2- State information service. (2020). Egypt's COVID-19 co-existence plan. Retrieved from <https://www.sis.gov.eg/Story/146118/-Egypt%E2%80%99s-COVID-19-co-existence-plan?lang=en-us>.
- 3- Mujika, I., & Padilla, S. (2000). Detraining: loss of training-induced physiological and performance adaptations. Part II: Long term insufficient training stimulus. *Sports medicine* (Auckland, N.Z.), 30(3), 145–154. <https://doi.org/10.2165/00007256-200030030-00001>
- 4- Chen, P., Mao, L., Nassis, G. P., Harmer, P., Ainsworth, B. E., & Li, F. (2020). Coronavirus disease (COVID-19): The need to maintain regular physical activity while taking precautions. *Journal of sport and health science*, 9(2), 103–104. <https://doi.org/10.1016/j.jshs.2020.02.001>
- 5- Joo C. H. (2018). The effects of short term detraining and retraining on physical fitness in elite soccer players. *PloS one*, 13(5), e0196212. <https://doi.org/10.1371/journal.pone.0196212>
- 6- Yoon J. (2002). Physiological profiles of elite senior wrestlers. *Sports medicine* (Auckland, N.Z.), 32(4), 225–233. <https://doi.org/10.2165/00007256-200232040-00002>
- 7- Arslanoğlu, E. (2015). Physical profiles of Turkish young Greco-Roman wrestlers. *Educational Research and Reviews*, 10(8), 1034 - 1038.
- 8- Ramirez-Velez, R., Argothy, R., Meneses-Echavez, J. F., Beatriz Sanchez-Puccini, M., Lopez-Alban, C. A., & Cohen, D. D. (2014). Anthropometric characteristics and physical performance of colombian elite male wrestlers. *Asian journal of sports medicine*, 5(4), e23810. <https://doi.org/10.5812/asjms.23810>
- 9- Hadžović-Džuvo, A., Valjevac, A., Lepara, O., Pjanić, S., Hadžimuratović, A., & Mekić, A. (2014). Oxidative stress status in elite athletes engaged in different sport disciplines. *Bosnian journal of basic medical sciences*, 14(2), 56–62. <https://doi.org/10.17305/bjbms.2014.2262>
- 10- Scribbans, T. D., Vecsey, S., Hankinson, P. B., Foster, W. S., & Gurd, B. J. (2016). The Effect of Training Intensity on VO₂max in Young Healthy Adults: A Meta-Regression and Meta-Analysis. *International journal of exercise science*, 9(2), 230–247.
- 11- Dlugosz, E. M., Chappell, M. A., Meek, T. H., Szafrńska, P. A., Zub, K., Konarzewski, M., et al. (2013). Phylogenetic analysis of mammalian maximal oxygen consumption during exercise. *Journal of Experimental Biology*, 216(24), 4712–4721. doi:10.1242/jeb.088914
- 12- Hübner-Woźniak, E., Kosmol, A., & Gajewski, J. (2009). Aerobic fitness of elite female and male wrestlers. *Biology of Sport*, 26(4), 339–348. <https://doi.org/10.5604/20831862.901138>
- 13- Heywood V. (2006). In: Advanced fitness assessment and exercise prescription. 5 ed. Champaign, IL: Human Kinetics; The physical fitness specialist manual, The Cooper institute for aerobics research, Dallas TX, revised 2005. Heywood V, editor.
- 14- Gulam A. (2016). Impact of exercises on muscular, respiratory and circulatory body systems. *IJAR*. 1;12:59-62
- 15- Nicolò, A., Massaroni, C., & Passfield, L. (2017). Respiratory Frequency during Exercise: The Neglected Physiological Measure. *Frontiers in physiology*, 8, 922. <https://doi.org/10.3389/fphys.2017.00922>
- 16- Ramos-Jiménez, A., Hernández-Torres, R. P., Torres-Durán, P. V., Romero-Gonzalez, J., Mascher, D., Posadas-Romero, C., & Juárez-Oropeza, M. A. (2008). The Respiratory Exchange Ratio is Associated with Fitness Indicators Both in Trained and Untrained Men: A Possible

- Application for People with Reduced Exercise Tolerance. *Clinical medicine. Circulatory, respiratory and pulmonary medicine*, 2, 1–9. <https://doi.org/10.4137/ccrpm.s449>
- 17- Bellar, D., & Judge, L. W. (2012). Modeling and relationship of respiratory exchange ratio to athletic performance. *Journal of strength and conditioning research*, 26(9), 2484–2489. <https://doi.org/10.1519/JSC.0b013e31823f271d>
- 18- Bearden, S. E., Henning, P. C., Bearden, T. A., & Moffatt, R. J. (2004). The slow component of VO(2) kinetics in very heavy and fatiguing square-wave exercise. *European journal of applied physiology*, 91(5-6), 586–594. <https://doi.org/10.1007/s00421-003-1009-x>
- 19- Houmard, J. A., Hortobágyi, T., Johns, R. A., Bruno, N. J., Nute, C. C., Shinebarger, M. H., & Welborn, J. W. (1992). Effect of short-term training cessation on performance measures in distance runners. *International journal of sports medicine*, 13(8), 572–576. <https://doi.org/10.1055/s-2007-1024567>
- 20- Rietjens, G. J., Keizer, H. A., Kuipers, H., & Saris, W. H. (2001). A reduction in training volume and intensity for 21 days does not impair performance in cyclists. *British journal of sports medicine*, 35(6), 431–434. <https://doi.org/10.1136/bjsm.35.6.431>
- 21- Madsen, K., Pedersen, P. K., Djurhuus, M. S., & Klitgaard, N. A. (1993). Effects of detraining on endurance capacity and metabolic changes during prolonged exhaustive exercise. *Journal of applied physiology (Bethesda, Md. : 1985)*, 75(4), 1444–1451. <https://doi.org/10.1152/jappl.1993.75.4.1444>
- 22- Mujika, I., & Padilla, S. (2001). Muscular characteristics of detraining in humans. *Medicine and science in sports and exercise*, 33(8), 1297–1303. <https://doi.org/10.1097/00005768-200108000-00009>
- 23- MacDougall JD, Sale D. (n.d.). The physiology of training for high performance. Retrieved from <https://www.amazon.com/Physiology-Training-High-Performance/dp/0199650640>
- 24- García-Pallarés, J., López-Gullón, J. M., Muriel, X., Díaz, A., & Izquierdo, M. (2011). Physical fitness factors to predict male Olympic wrestling performance. *European journal of applied physiology*, 111(8), 1747–1758. <https://doi.org/10.1007/s00421-010-1809-8>
- 25- Iermakov, S., Podrigalo, L., & Jagiełło, W. (2016). Hand-grip strength as an indicator for predicting the success in martial arts athletes. *Archives of Budo*, 12.
- 26- Gentil, P., Soares, S., & Bottaro, M. (2015). Single vs. Multi-Joint Resistance Exercises: Effects on Muscle Strength and Hypertrophy. *Asian journal of sports medicine*, 6(2), e24057. <https://doi.org/10.5812/asjsm.24057>
- 27- Guilhem, G., Giroux, C., Couturier, A., & Maffiuletti, N. A. (2014). Validity of trunk extensor and flexor torque measurements using isokinetic dynamometry. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology*, 24(6), 986–993. <https://doi.org/10.1016/j.jelekin.2014.07.006>
- 28- Kraemer WJ, Fry AC, Rubin MR, Triplett-McBride T, Gordon SE, Koziris LP, Lynch JM, Volek JS, Meuffels DE, Newton RU, Fleck SJ. Physiological and performance responses to tournament wrestling. *Med Sci Sports Exerc*. 2001 Aug;33(8):1367-78. doi: 10.1097/00005768-200108000-00019. PMID: 11474340.
- 29- de França, H. S., Branco, P. A., Guedes Junior, D. P., Gentil, P., Steele, J., & Teixeira, C. V. (2015). The effects of adding single-joint exercises to a multi-joint exercise resistance training program on upper body muscle strength and size in trained men. *Applied physiology, nutrition,*

and metabolism = *Physiologie appliquee, nutrition et metabolisme*, 40(8), 822–826.
<https://doi.org/10.1139/apnm-2015-0109>

30- Tucci, J. T., Carpenter, D. M., Pollock, M. L., Graves, J. E., & Leggett, S. H. (1992). Effect of reduced frequency of training and detraining on lumbar extension strength. *Spine*, 17(12), 1497–1501. <https://doi.org/10.1097/00007632-199212000-00009>

31- Basar, S., Duzgun, I., Guzel, N. A., Cicioğlu, I., & Celik, B. (2014). Differences in strength, flexibility and stability in freestyle and Greco-Roman wrestlers. *Journal of back and musculoskeletal rehabilitation*, 27(3), 321–330. <https://doi.org/10.3233/BMR-130451>

32- Chaabene, H., Negra, Y., Bouguezzi, R., Mkaouer, B., Franchini, E., Julio, U., & Hachana, Y. (2017). Physical and Physiological Attributes of Wrestlers: An Update. *Journal of strength and conditioning research*, 31(5), 1411–1442. <https://doi.org/10.1519/JSC.0000000000001738>

33- Leite, T. B., Costa, P. B., Leite, R. D., Novaes, J. S., Fleck, S. J., & Simão, R. (2017). Effects of Different Number of Sets of Resistance Training on Flexibility. *International journal of exercise science*, 10(3), 354–364.

34- Saraiva, A. R., Reis, V. M., Costa, P. B., Bentes, C. M., Costa E Silva, G. V., & Novaes, J. S. (2014). Chronic effects of different resistance training exercise orders on flexibility in elite judo athletes. *Journal of human kinetics*, 40, 129–137. <https://doi.org/10.2478/hukin-2014-0015>

35- Swärd, L., Svensson, M., & Zetterberg, C. (1990). Isometric muscle strength and quantitative electromyography of back muscles in wrestlers and tennis players. *The American journal of sports medicine*, 18(4), 382–386. <https://doi.org/10.1177/036354659001800409>